

EXERCISE - 7.1

1. In quadrilateral $ACBD$, we have $AC = AD$ and AB being the bisector of $\angle A$.

Now, in $\triangle ABC$ and $\triangle ABD$,

$$AC = AD \quad \text{[Given]}$$

$$\angle CAB = \angle DAB \quad \text{[}\because AB \text{ bisects } \angle CAD\text{]}$$

$$AB = AB \quad \text{[Common]}$$

$$\therefore \triangle ABC \cong \triangle ABD \quad \text{[By SAS congruence]}$$

$$\text{So, } BC = BD \quad \text{[By C.P.C.T.]}$$

2. In quadrilateral $ABCD$, we have

$$AD = BC \text{ and } \angle DAB = \angle CBA$$

In $\triangle ABD$ and $\triangle BAC$,

$$AD = BC \quad \text{[Given]}$$

$$\angle DAB = \angle CBA \quad \text{[Given]}$$

$$AB = AB \quad \text{[Common]}$$

$$\therefore \triangle ABD \cong \triangle BAC \quad \text{[By SAS congruence]}$$

$$\therefore BD = AC \quad \text{[By C.P.C.T.]}$$

$$\text{and } \angle ABD = \angle BAC \quad \text{[By C.P.C.T.]}$$

3. In $\triangle OBC$ and $\triangle OAD$, we have

$$\angle OBC = \angle OAD \quad \text{[Each equals } 90^\circ\text{]}$$

$$BC = AD \quad \text{[Given]}$$

$$\angle BOC = \angle AOD \quad \text{[Vertically opposite angles]}$$

$$\therefore \triangle OBC \cong \triangle OAD \quad \text{[By AAS congruence]}$$

$$\Rightarrow OB = OA \quad \text{[By C.P.C.T.]}$$

i.e., O is the mid-point of AB .

Thus, CD bisects AB .

4. $\because p \parallel q$ and AC is a transversal.

$$\therefore \angle BAC = \angle DCA \quad \dots\text{(i)}$$

[Alternate interior angles]

Also $l \parallel m$ and AC is a transversal.

$$\therefore \angle BCA = \angle DAC \quad \dots\text{(ii)}$$

[Alternate interior angles]

Now, in $\triangle ABC$ and $\triangle CDA$, we have

$$\angle BAC = \angle DCA \quad \text{[From (i)]}$$

$$AC = AC \quad \text{[Common]}$$

$$\angle BCA = \angle DAC \quad \text{[From (ii)]}$$

$$\therefore \triangle ABC \cong \triangle CDA \quad \text{[By ASA congruence]}$$

5. We have, l is the bisector of $\angle QAP$.

$$\therefore \angle QAB = \angle PAB \text{ and}$$

$$\angle Q = \angle P \quad \text{[Each equals } 90^\circ\text{]}$$

$$\Rightarrow \angle ABQ = \angle ABP \quad \dots\text{(i)}$$

[By angle sum property of triangle]

(i) Now, in $\triangle APB$ and $\triangle AQB$, we have

$$\angle ABP = \angle ABQ \quad \text{[From (i)]}$$

$$AB = AB \quad \text{[Common]}$$

$$\angle PAB = \angle QAB \quad \text{[Given]}$$

$$\therefore \triangle APB \cong \triangle AQB \quad \text{[By ASA congruence]}$$

(ii) Since $\triangle APB \cong \triangle AQB$

$$\Rightarrow BP = BQ \quad \text{[By C.P.C.T.]}$$

or Perpendicular distance of B from AP
= Perpendicular distance of B from AQ

Thus, the point B is equidistant from the arms of $\angle A$.

6. We have, $\angle BAD = \angle EAC$

Adding $\angle DAC$ on both sides, we have

$$\angle BAD + \angle DAC = \angle EAC + \angle DAC$$

$$\Rightarrow \angle BAC = \angle DAE$$

Now, in $\triangle ABC$ and $\triangle ADE$, we have

$$\angle BAC = \angle DAE \quad \text{[Proved above]}$$

$$AB = AD \quad \text{[Given]}$$

$$AC = AE \quad \text{[Given]}$$

$$\therefore \triangle ABC \cong \triangle ADE \quad \text{[By SAS congruence]}$$

$$\Rightarrow BC = DE \quad \text{[By C.P.C.T.]}$$

7. We have, P is the mid-point of AB .

$$\therefore AP = BP$$

$$\angle EPA = \angle DPB \quad \text{[Given]}$$

Adding $\angle EPD$ on both sides, we get

$$\angle EPA + \angle EPD = \angle DPB + \angle EPD$$

$$\Rightarrow \angle APD = \angle BPE$$

(i) Now, in $\triangle DAP$ and $\triangle EBP$, we have

$$\angle PAD = \angle PBE \quad \text{[}\because \angle BAD = \angle ABE\text{]}$$

$$AP = BP \quad \text{[Proved above]}$$

$$\angle DPA = \angle EPB \quad \text{[Proved above]}$$

$$\therefore \triangle DAP \cong \triangle EBP \quad \text{[By ASA congruence]}$$

(ii) Since, $\triangle DAP \cong \triangle EBP$

$$\Rightarrow AD = BE \quad \text{[By C.P.C.T.]}$$

8. Since M is the mid-point of AB .

$$\therefore BM = AM \quad \text{[Given]}$$

(i) In $\triangle AMC$ and $\triangle BMD$, we have

$$CM = DM \quad \text{[Given]}$$

$$\angle AMC = \angle BMD \quad \text{[Vertically opposite angles]}$$

$$AM = BM \quad \text{[Proved above]}$$

$$\therefore \triangle AMC \cong \triangle BMD \quad \text{[By SAS congruence]}$$

(ii) Since $\triangle AMC \cong \triangle BMD$

$$\Rightarrow \angle MAC = \angle MBD \quad \text{[By C.P.C.T.]}$$

But they form a pair of alternate interior angles.

$$\therefore AC \parallel DB$$

Now, BC is a transversal which intersects parallel lines AC and DB .

$$\therefore \angle BCA + \angle DBC = 180^\circ \quad [\text{Co-interior angles}]$$

$$\text{But } \angle BCA = 90^\circ \quad [\because \triangle ABC \text{ is right angled at } C]$$

$$\therefore 90^\circ + \angle DBC = 180^\circ$$

$$\Rightarrow \angle DBC = 90^\circ$$

(iii) Since, $\triangle AMC \cong \triangle BMD$

$$\therefore AC = BD \quad [\text{By C.P.C.T.}]$$

Now, in $\triangle DBC$ and $\triangle ACB$, we have

$$BD = CA \quad [\text{Proved above}]$$

$$\angle DBC = \angle ACB \quad [\text{Each equals } 90^\circ]$$

$$BC = CB \quad [\text{Common}]$$

$$\therefore \triangle DBC \cong \triangle ACB \quad [\text{By SAS congruence}]$$

(iv) As $\triangle DBC \cong \triangle ACB$

$$\Rightarrow DC = AB \quad [\text{By C.P.C.T.}]$$

$$\text{But } DM = CM \quad [\text{Given}]$$

$$\therefore CM = \frac{1}{2}DC = \frac{1}{2}AB$$

EXERCISE - 7.2

1. (i) Let BD and CE are the bisectors of $\angle B$ and $\angle C$ respectively.

In $\triangle ABC$, we have $AC = AB$

$$\therefore \angle ABC = \angle ACB$$

[Angles opposite to equal sides of a triangle are equal]

$$\Rightarrow \frac{1}{2}\angle ABC = \frac{1}{2}\angle ACB$$

$$\Rightarrow \angle OBC = \angle OCB$$

$$\Rightarrow OC = OB$$

[Sides opposite to equal angles of a triangle are equal]

(ii) In $\triangle ABO$ and $\triangle ACO$, we have

$$AB = AC \quad [\text{Given}]$$

$$\angle OBA = \angle OCA$$

$$\left[\because \frac{1}{2}\angle B = \frac{1}{2}\angle C \right]$$

$$OB = OC \quad [\text{Proved above}]$$

$$\therefore \triangle ABO \cong \triangle ACO \quad [\text{By SAS congruence}]$$

$$\Rightarrow \angle OAB = \angle OAC \quad [\text{By C.P.C.T.}]$$

$$\Rightarrow AO \text{ bisects } \angle A.$$

2. Since AD is bisector of BC .

$$\therefore BD = CD$$

Now, in $\triangle ABD$ and $\triangle ACD$, we have

$$AD = AD \quad [\text{Common}]$$

$$\angle ADB = \angle ADC \quad [\text{Each equals } 90^\circ]$$

$$BD = CD \quad [\text{Proved above}]$$

$$\therefore \triangle ABD \cong \triangle ACD \quad [\text{By SAS congruence}]$$

$$\Rightarrow AB = AC \quad [\text{By C.P.C.T.}]$$

Thus, ABC is an isosceles triangle.

3. Given, $\triangle ABC$ is an isosceles triangle with $AB = AC$

$$\Rightarrow \angle ACB = \angle ABC$$

[Angles opposite to equal sides of a triangle are equal]

$$\Rightarrow \angle BCE = \angle CBF$$

Now, in $\triangle BEC$ and $\triangle CFB$, we have

$$\angle BCE = \angle CBF \quad [\text{Proved above}]$$

$$\angle BEC = \angle CFB \quad [\text{Each equals } 90^\circ]$$

$$BC = CB \quad [\text{Common}]$$

$$\therefore \triangle BEC \cong \triangle CFB \quad [\text{By AAS congruence}]$$

$$\text{So, } BE = CF \quad [\text{By C.P.C.T.}]$$

4. (i) In $\triangle ABE$ and $\triangle ACF$, we have

$$\angle AEB = \angle AFC \quad [\text{Each } 90^\circ \text{ as } BE \perp AC \text{ and } CF \perp AB]$$

$$\angle A = \angle A \quad [\text{Common}]$$

$$BE = CF \quad [\text{Given}]$$

$$\therefore \triangle ABE \cong \triangle ACF \quad [\text{By AAS congruence}]$$

(ii) Since, $\triangle ABE \cong \triangle ACF$

$$\therefore AB = AC \quad [\text{By C.P.C.T.}]$$

5. In $\triangle ABC$, we have

$$AB = AC \quad [\because ABC \text{ is an isosceles triangle}]$$

$$\therefore \angle ABC = \angle ACB \quad \dots(i)$$

[Angles opposite to equal sides of a triangle are equal]

Again, in $\triangle BDC$, we have

$$BD = CD \quad [\because BDC \text{ is an isosceles triangle}]$$

$$\therefore \angle CBD = \angle BCD \quad \dots(ii)$$

[Angles opposite to equal sides of a triangle are equal]

Adding (i) and (ii), we have

$$\angle ABC + \angle CBD = \angle ACB + \angle BCD$$

$$\Rightarrow \angle ABD = \angle ACD$$

6. Given, $AB = AC$ and $AB = AD$

$$\therefore AC = AD$$

Now, in $\triangle ABC$, we have

$$\angle B + \angle ACB + \angle BAC = 180^\circ \quad [\text{Angle sum property of a triangle}]$$

$$\Rightarrow 2\angle ACB + \angle BAC = 180^\circ \quad \dots(i)$$

$$[\because \angle B = \angle ACB \text{ (Angles opposite to equal sides of a triangle are equal)}]$$

Similarly, in $\triangle ACD$,

$$\angle D + \angle ACD + \angle CAD = 180^\circ \quad [\text{Angle sum property of a triangle}]$$

$$\Rightarrow 2\angle ACD + \angle CAD = 180^\circ \quad \dots(ii)$$

$$[\because \angle D = \angle ACD \text{ (Angles opposite to equal sides of a triangle are equal)}]$$

Adding (i) and (ii), we have

$$2\angle ACB + \angle BAC + 2\angle ACD + \angle CAD = 180^\circ + 180^\circ$$

$$\Rightarrow 2[\angle ACB + \angle ACD] + [\angle BAC + \angle CAD] = 360^\circ$$

$$\Rightarrow 2\angle BCD + 180^\circ = 360^\circ$$

$$[\because \angle BAC \text{ and } \angle CAD \text{ form a linear pair}]$$

$$\Rightarrow 2\angle BCD = 360^\circ - 180^\circ = 180^\circ$$

$$\Rightarrow \angle BCD = \frac{180^\circ}{2} = 90^\circ$$

7. In $\triangle ABC$, we have

$$AB = AC \quad [\text{Given}]$$

$$\therefore \angle ACB = \angle ABC$$

[Angles opposite to equal sides of a triangle are equal]

$$\text{Now, } \angle A + \angle B + \angle C = 180^\circ \quad [\text{Angle sum property of a triangle}]$$



$$\Rightarrow 90^\circ + \angle B + \angle C = 180^\circ \quad [\angle A = 90^\circ (\text{Given})]$$

$$\Rightarrow \angle B + \angle C = 180^\circ - 90^\circ = 90^\circ$$

$$\text{But } \angle ABC = \angle ACB, \text{ i.e., } \angle B = \angle C$$

$$\therefore \angle B = \angle C = \frac{90^\circ}{2} = 45^\circ$$

$$\text{Thus, } \angle B = 45^\circ \text{ and } \angle C = 45^\circ$$



8. In $\triangle ABC$, we have

$AB = BC = CA$ [$\because ABC$ is an equilateral triangle]

Now, $AB = BC \Rightarrow \angle A = \angle C$... (i)

[Angles opposite to equal sides of a triangle are equal]

Similarly, $AC = BC \Rightarrow \angle A = \angle B$... (ii)

From (i) and (ii), we have $\angle A = \angle B = \angle C$

Let $\angle A = \angle B = \angle C = x$

Since, $\angle A + \angle B + \angle C = 180^\circ$


[Angle sum property of a triangle]

$\therefore x + x + x = 180^\circ \Rightarrow 3x = 180^\circ$

$\Rightarrow x = 60^\circ$

$\therefore \angle A = \angle B = \angle C = 60^\circ$

Thus, the angles of an equilateral triangle are 60° each.



EXERCISE - 7.3

1. (i) In $\triangle ABD$ and $\triangle ACD$, we have

$AB = AC$ [Given]

$AD = AD$ [Common]

$BD = CD$ [Given]

$\therefore \triangle ABD \cong \triangle ACD$ [By SSS congruence]

$\Rightarrow \angle BAD = \angle CAD$ [By C.P.C.T.]

$\Rightarrow \angle BAP = \angle CAP$... (1)

(ii) In $\triangle ABP$ and $\triangle ACP$, we have

$AB = AC$ [Given]

$\angle BAP = \angle CAP$ [From (1)]

$AP = AP$ [Common]

$\therefore \triangle ABP \cong \triangle ACP$ [By SAS congruence]

(iii) Since, $\angle BAP = \angle CAP$ [From (1)]

$\therefore AP$ is the bisector of $\angle A$.

Again, in $\triangle BDP$ and $\triangle CDP$, we have

$BD = CD$ [Given]

$DP = DP$ [Common]

$BP = CP$ [$\because \triangle ABP \cong \triangle ACP$]

$\therefore \triangle BDP \cong \triangle CDP$ [By SSS congruence]

$\Rightarrow \angle BDP = \angle CDP$ [By C.P.C.T.]

$\Rightarrow DP$ (or AP) is the bisector of $\angle BDC$

$\therefore AP$ is the bisector of $\angle A$ as well as $\angle D$.

(iv) As, $\triangle ABP \cong \triangle ACP$

$\Rightarrow \angle APB = \angle APC$ and $BP = CP$ [By C.P.C.T.]

But $\angle APB + \angle APC = 180^\circ$ [Linear pair]

$\therefore \angle APB = \angle APC = 90^\circ$

$\Rightarrow AP \perp BC$.

Hence, AP is the perpendicular bisector of BC .

2. (i) In right $\triangle ABD$ and $\triangle ACD$,

$AB = AC$ [Given]

$\angle ADB = \angle ADC$ [Each equals 90°]

$AD = AD$ [Common]

$\therefore \triangle ABD \cong \triangle ACD$ [By RHS congruence]


So, $BD = CD$ [By C.P.C.T.]

$\Rightarrow D$ is the mid-point of BC or AD bisects BC .

(ii) Since, $\triangle ABD \cong \triangle ACD$,

$\Rightarrow \angle BAD = \angle CAD$ [By C.P.C.T.]

So, AD bisects $\angle A$.



3. In $\triangle ABC$, AM is the median [Given]

$\therefore BM = \frac{1}{2}BC$... (1)

In $\triangle PQR$, PN is the median [Given]

$\therefore QN = \frac{1}{2}QR$... (2)

Also, $BC = QR \Rightarrow \frac{1}{2}BC = \frac{1}{2}QR$

$\Rightarrow BM = QN$ [From (1) and (2)] ... (3)

(i) In $\triangle ABM$ and $\triangle PQN$, we have

$AB = PQ$ [Given]

$AM = PN$ [Given]

$BM = QN$ [From (3)]

$\therefore \triangle ABM \cong \triangle PQN$ [By SSS congruence]

(ii) Since $\triangle ABM \cong \triangle PQN$

$\Rightarrow \angle B = \angle Q$ [By C.P.C.T.] ... (4)

Now, in $\triangle ABC$ and $\triangle PQR$, we have

$\angle B = \angle Q$ [From (4)]

$AB = PQ$ [Given]

$BC = QR$ [Given]

$\therefore \triangle ABC \cong \triangle PQR$ [By SAS congruence]

4. Since, $BE \perp AC$ [Given]

$\therefore \triangle BEC$ is a right triangle such that

$\angle BEC = 90^\circ$

Similarly, $\angle CFB = 90^\circ$

Now, in right $\triangle BEC$ and right $\triangle CFB$,

we have

$BE = CF$

$BC = CB$

$\angle BEC = \angle CFB = 90^\circ$

$\therefore \triangle BEC \cong \triangle CFB$

[By RHS congruence]

So, $\angle BCE = \angle CBF$

[By C.P.C.T.]

or $\angle BCA = \angle CBA$

Now, in $\triangle ABC$, $\angle BCA = \angle CBA \Rightarrow AB = AC$

[Sides opposite to equal angles of a triangle are equal]

$\therefore \triangle ABC$ is an isosceles triangle.

5. We have, $AP \perp BC$

$\therefore \angle APB = 90^\circ$ and $\angle APC = 90^\circ$

In $\triangle ABP$ and $\triangle ACP$, we have


$\angle APB = \angle APC$ [Each equals 90°]

$AB = AC$ [Given]

$AP = AP$ [Common]


$\therefore \triangle ABP \cong \triangle ACP$ [By RHS congruence]

So, $\angle B = \angle C$ [By C.P.C.T.]



[Given]

[Common]




EXERCISE - 7.5

1. Let us consider a $\triangle ABC$.

Draw l , the perpendicular bisector of AB .

Draw m , the perpendicular bisector of BC .

Let the two perpendicular bisectors l and m meet at O .



O is the required point which is equidistant from all the vertices A , B and C .

Note: If we draw a circle with centre O and radius OA , OB or OC , then it will pass through A , B and C . The point O is called circumcentre of the triangle.

2. Let us consider a $\triangle ABC$.

Draw l , the bisector of $\angle B$.



Draw m , the bisector of $\angle C$.

Let the two bisectors l and m meet at O .

Thus, O is the required point which is equidistant from the sides of $\triangle ABC$.

Note: If we draw $OM \perp BC$ and draw a circle with O as centre and OM as radius, then the circle will touch the sides of the triangle.

Point O is called the incentre of the triangle.



3. Let us join A and B , and draw l , the perpendicular bisector of AB .

Now, join B and C , and draw m , the perpendicular bisector of BC . Let the perpendicular bisectors l and m meet at O . The point O is the required point where the ice cream parlour be set up.

Note: If we join A and C and draw the perpendicular bisectors, then it will also meet (or pass through) the point O .

4. It is an activity. We require 150 equilateral triangles of side 1 cm in the figure (i) and 300 equilateral triangles in the figure (ii).

\therefore The figure (ii) has more triangles.

