# **Probability**



### **SOLUTIONS**

#### EXERCISE - 15.1

- **1.** (i) Probability of an event E + Probability of the event 'not E' = 1.
- (ii) The probability of an event that cannot happen is 0. Such an event is called impossible event.
- (iii) The probability of an event that is certain to happen is 1. Such an event is called sure or certain event.
- (iv) The sum of the probabilities of all the elementary events of an experiment is 1.
- (v) The probability of an event is greater than or equal to 0 and less than or equal to 1.
- **2.** (i) Since the driver may or may not start the car, thus the outcomes are not equally likely.
- (ii) The player may shoot or miss the shot.
- :. The outcomes are not equally likely.
- (iii) In advance it is known that the answer is to be either right or wrong.
- ... The outcomes right or wrong are equally likely to occur.
- (iv) In advance it is known that newly born baby has to be either a boy or a girl.
- ... The outcomes either a boy or a girl are equally likely to occur.
- 3. Since on tossing a coin, the outcomes 'head' and 'tail' are equally likely, the result of tossing a coin is completely unpredictable and so it is a fair way.
- **4. (b)** : Since, the probability of an event cannot be negative.
- ∴ -1.5 cannot be the probability of an event.
- 5. : P(E) + P(not E) = 1
- ..  $0.05 + P(\text{not } E) = 1 \Rightarrow P(\text{not } E) = 1 0.05 = 0.95$ Thus, probability of 'not E' = 0.95.
- **6.** (i) Since there are lemon flavoured candies only in the bag.
- :. Taking out orange flavoured candy is not possible.
- $\Rightarrow$  Probability of taking out an orange flavoured candy = 0.
- (ii) Probability of taking out a lemon flavoured candy = 1.
- **7.** Let the probability of 2 students having same birthday = P(SB)

And the probability of 2 students not having the same birthday = P(NSB)

- $\therefore P(SB) + P(NSB) = 1$
- $\Rightarrow$   $P(SB) + 0.992 = 1 \Rightarrow P(SB) = 1 0.992 = 0.008$

- 8. Total number of balls = 3 + 5 = 8
- :. Number of possible outcomes = 8
- (i) ∴ There are 3 red balls.
- $\Rightarrow$  Number of favourable outcomes = 3
- $\therefore P(\text{red ball}) = \frac{\text{Number of favourable outcomes}}{\text{Number of all possible outcomes}} = \frac{3}{8}$
- (ii) Probability of the ball drawn which is not red

$$= 1 - P(\text{red ball}) = 1 - \frac{3}{8} = \frac{8 - 3}{8} = \frac{5}{8}$$

- 9. Total number of marbles = 5 + 8 + 4 = 17
- :. Number of all possible outcomes = 17
- (i) : Number of red marbles = 5
- ⇒ Number of favourable outcomes = 5
- $\therefore$  Probability of red marbles,  $P(\text{red}) = \frac{5}{17}$
- (ii) : Number of white marbles = 8
- $\Rightarrow$  Number of favourable outcomes = 8
- $\therefore$  Probability of white marbles,  $P(\text{white}) = \frac{8}{17}$
- (iii) ∴ Number of green marbles = 4
- $\therefore$  Number of marbles which are not green = 17 4 = 13
- ⇒ Number of favourable outcomes = 13
- : Probability of marbles 'not green',  $P(\text{not green}) = \frac{13}{17}$
- **10.** Number of : 50 p coins = 100, ₹ 1 coins = 50 ₹ 2 coins = 20, ₹ 5 coins = 10

Total number of coins = 100 + 50 + 20 + 10 = 180

- :. Total possible outcomes = 180
- (i) Number of favourable outcomes = 100

$$\therefore$$
  $P(50 \text{ p coins}) = \frac{100}{180} = \frac{5}{9}$ 

- (ii) Number of ₹5 coins = 10
- $\therefore$  Number of 'not ₹ 5' coins = 180 10 = 170
- $\Rightarrow$  Number of favourable outcomes = 170
- ∴  $P(\text{not } ₹ 5 \text{ coin}) = \frac{170}{180} = \frac{17}{18}$
- 11. Number of male fishes = 5
- Number of female fishes = 8 ∴ Total number of fishes = 5 + 8 = 13
- $\Rightarrow$  Total number of outcomes = 13
- $\therefore$  *P*(fish taken out is a male fish) = 5/13.
- **12.** Total number marked = 8
- :. Total number of possible outcomes = 8
- (i) Number of favourable outcomes = 1
- $P(8) = \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{1}{8}$

- (ii) Odd numbers are 1, 3, 5 and 7.
- $\therefore$  Number of odd numbers from 1 to 8 = 4
- ⇒ Number of favourable outcomes = 4
- $\therefore$  *P* (an odd number)
  - $= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{4}{8} = \frac{1}{2}$
- (iii) The numbers 3, 4, 5, 6, 7 and 8 are greater than 2.
- $\therefore$  Number of numbers greater than 2 = 6
- $\Rightarrow$  Number of favourable outcomes = 6
- $\therefore$  P(a number greater than 2)

$$= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{6}{8} = \frac{3}{4}$$

- (iv) The numbers 1, 2, 3, 4, 5, 6, 7 and 8 are less than 9.
- $\therefore$  Number of numbers less than 9 = 8
- $\Rightarrow$  Number of favourable outcomes = 8
- $\therefore$  *P*(a number less than 9)

$$= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{8}{8} = 1$$

- **13.** Since, numbers on a die are 1, 2, 3, 4, 5 and 6.
- :. Number of total possible outcomes = 6
- (i) Since 2, 3 and 5 are prime numbers.
- $\Rightarrow$  Number of favourable outcomes = 3 P(a prime number)

$$= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{3}{6} = \frac{1}{2}$$

- (ii) Since, the numbers between 2 and 6 are 3, 4 and 5.
- $\Rightarrow$  Number of favourable outcomes = 3
- ∴ *P*(a number lying between 2 and 6)

$$= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{3}{6} = \frac{1}{2}$$

- (iii) Since 1, 3 and 5 are odd numbers.
- $\Rightarrow$  Number of favourable outcomes = 3
- $\therefore$  P(an odd number)

$$= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{3}{6} = \frac{1}{2}$$

- 14. Number of cards in deck = 52
- : Total number of possible outcomes = 52
- (i) ∴ Number of red colour kings = 2

[: King of diamond and heart is red]

- $\Rightarrow$  Number of favourable outcomes = 2
- ∴  $P(\text{a red king}) = \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}}$ =  $\frac{2}{52} = \frac{1}{26}$
- (ii) : 4 kings, 4 queens and 4 jacks are face cards.
- $\therefore$  Number of face cards = 12
- $\Rightarrow$  Number of favourable outcomes = 12
- ∴  $P(\text{a face card}) = \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}}$  $= \frac{12}{52} = \frac{3}{13}$
- (iii) Since, cards of diamond and heart are red.
- ... There are 2 kings, 2 queens, 2 jacks *i.e.*, 6 cards are red face cards.

- $\Rightarrow$  Number of favourable outcomes = 6
  - P(red face card) =  $\frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}}$

$$=\frac{6}{52}=\frac{3}{26}$$

- (iv) Since, there is only 1 jack of hearts.
- $\Rightarrow$  Number of favourable outcomes = 1
- $\therefore$  *P*(jack of hearts)

$$= \frac{Number of favourable outcomes}{Total number of possible outcomes} = \frac{1}{52}$$

- (v) ∵ There are 13 spades in a pack of 52 cards.
- $\Rightarrow$  Number of favourable outcomes = 13
- $P(a \text{ spade}) = \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}}$

$$=\frac{13}{52}=\frac{1}{4}$$

- (vi) ∵ There is only one queen of diamonds.
- $\Rightarrow$  Number of favourable outcomes = 1
- ∴ P(a queen of diamonds)

$$= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{1}{52}$$

- 15. We have five cards.
- ∴ Total number of possible outcomes = 5
- (i) ∵ Number of queens = 1
- $\Rightarrow$  Number of favourable outcomes = 1

$$\therefore P(\text{a queen}) = \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{1}{5}$$

- (ii) The queen is drawn and put aside.
- $\Rightarrow$  Only 5 1 = 4 cards are left.
- : Total number of possible outcomes = 4
- (a) ∵ There is only one ace.
- $\Rightarrow$  Number of favourable outcomes = 1

$$\therefore P(\text{an ace}) = \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{1}{4}$$

- (b) Since, the only queen has already been put aside.
- $\Rightarrow$  Number of possible outcomes = 0
- $\therefore$  P(a queen)

$$= \frac{Number of favourable outcomes}{Total number of possible outcomes} = \frac{0}{4} = 0$$

- **16.** We have, number of good pens = 132 and number of defective pens = 12
- $\Rightarrow$  Total number of possible outcomes = 132 + 12 = 144
- ⇒ Number of favourable outcomes = 132
- $\therefore$  P(good pens)

$$= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{132}{144} = \frac{11}{12}$$

- **17.** (i) Since, there are 20 bulbs in the lot.
- $\Rightarrow$  Total number of possible outcomes = 20
- : Number of defective bulbs = 4
- $\Rightarrow$  Number of favourable outcomes = 4
- ∴ *P*(defective bulb)
  - $= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{4}{20} = \frac{1}{5}$

Probability 3

- (ii) : The bulb drawn above is not included in the lot.
- Number of remaining bulbs = 20 1 = 19
- Total number of possible outcomes = 19
- Number of bulbs which are not defective = 19 - 4 = 15
- Number of favourable outcomes = 15
- *P*(not defective bulb)
  - Number of favourable outcomes =  $\frac{15}{}$ Total number of possible outcomes
- We have, total number of discs = 90
- Total number of possible outcomes = 90
- (i) Since the two-digit numbers are 10, 11, 12, ..., 90.
- Number of two-digit numbers = 90 9 = 81
- Number of favourable outcomes = 81  $\Rightarrow$
- *P*(a two-digit number)
  - $\frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{81}{90} = \frac{9}{10}$
- (ii) Perfect square numbers from 1 to 90 are 1, 4, 9, 16, 25, 36, 49, 64 and 81.
- Number of perfect squares = 9∴.
- Number of favourable outcomes = 9
- *P*(a perfect square number)
  - $\frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{9}{90} = \frac{1}{10}$
- (iii) Numbers divisible by 5 from 1 to 90 are 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90
- i.e., There are 18 numbers from 1 to 90 which are divisible by 5.
- Number of favourable outcomes = 18  $\Rightarrow$
- *P*(a number divisible by 5)
  - $\frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{18}{90} = \frac{1}{5}$
- 19. Since there are six faces of the given die and these faces are marked with letters A, B, C, D, E and A.
- :. Total number of letters = 6
- $\Rightarrow$ Total number of possible outcomes = 6
- Two faces are having the letter A. (i)
- Number of favourable outcomes = 2
- *P*(getting letter A)
  - Number of favourable outcomes  $=\frac{2}{6}=\frac{1}{3}$ Total number of possible outcomes 6
- Only one face is having the letter D. (ii)
- Number of favourable outcomes = 1
- *P*(getting letter D)
  - $= \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}} = \frac{1}{6}$
- 20. Here, area of the rectangle =  $3 \text{ m} \times 2 \text{ m} = 6 \text{ m}^2$

And, the area of the circle  $= \pi r^2 = \pi \left(\frac{1}{2}\right)^2 m^2 = \frac{\pi}{4} m^2$ 

- Probability for the die to fall inside the circle
- Area of the favourable region

Area of the whole region

$$= \frac{\text{Area of the circle}}{\text{Area of the rectangle}} = \frac{\left[\frac{\pi}{4}\right]}{6} = \frac{\pi}{4} \times \frac{1}{6} = \frac{\pi}{24}$$

- Total number of ball pens = 14421.
- Total number of possible outcomes = 144  $\Rightarrow$
- (i) Since there are 20 defective pens.
- Number of good pens = 144 20 = 124
- Number of favourable outcomes = 124
- Probability that she will buy it =  $\frac{124}{144} = \frac{31}{36}$
- (ii) Probability that she will not buy it
- = 1 [Probability that she will buy it]

$$=1-\frac{31}{36}=\frac{36-31}{36}=\frac{5}{36}$$

- **22.** : The two dice are thrown together.
- Following are the possible outcomes:
- $\{(1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 3), (2, 1), (2, 2), (2, 3), (2, 1), (2, 2), (2, 3), (2, 3), (2, 4), (2,$
- (2, 4), (2, 5), (2, 6), (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6),
- (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), (5, 1), (5, 2), (5, 3),
- (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6).
- Total number of possible outcomes =  $6 \times 6 = 36$
- (a) : The sum on two dice is 3 for: (1, 2) and (2, 1)
- Number of favourable outcomes =  $2 \Rightarrow P(3) = 2/36$
- (p) :: The sum on two dice is 4 for: (1, 3), (2, 2) and (3, 1).
- Number of favourable outcomes =  $3 \Rightarrow P(4) = 3/36$
- (c) The sum on two dice is 5 for: (1, 4), (2, 3), (3, 2) and (4, 1)
- Number of favourable outcomes =  $4 \Rightarrow P(5) = 4/36$
- (d) :: The sum on two dice is 6 for: (1, 5), (2, 4), (3, 3), (4, 2) and (5, 1)
- Number of favourable outcomes =  $5 \Rightarrow P(6) = 5/36$
- (e) : The sum on two dice is 7 for : (1, 6), (2, 5), (3, 4), (4, 3), (5, 2) and (6, 1)
- Number of favourable outcomes =  $6 \Rightarrow P(7) = 6/36$
- ∴ The sum on two dice is 9 for : (3, 6), (4, 5), (5, 4) and (6, 3)
- Number of favourable outcomes =  $4 \Rightarrow P(9) = 4/36$
- (g) : The sum on two dice is 10 for: (4, 6), (5, 5) and (6, 4)
- Number of favourable outcomes =  $3 \Rightarrow P(10) = 3/36$
- $\therefore$  The sum on two dice is 11 for : (5, 6) and (6, 5)
- Number of favourable outcomes =  $2 \Rightarrow P(11) = 2/36$ Thus, the complete table is as under:

Event: 'Sum on 2 dice'	Probability		
2	1/36		
3	2/36		
4	3/36		
5	4/36		
6	5/36		
7	6/36		
8	5/36		
9	4/36		
10	3/36		
11	2/36		
12	1/36		

- (ii) No, the number of all possible outcomes is 36 not 11.
- :. The argument is not correct.
- **23.** All the possible outcomes are:

{HHH, HHT, HTT, TTT, TTH, THT, THH, HTH}

 $\therefore$  Number of all possible outcomes = 8

Let the event that Hanif will lose the game be denoted by *E*.

- ∴ Favourable outcomes are: {HHT, HTH, THH, THT, TTH, HTT}
- $\Rightarrow$  Number of favourable outcomes = 6
- $\therefore P(E) = \frac{6}{8} = \frac{3}{4}$
- **24.** Since, throwing a die twice or throwing two dice simultaneously is the same.
- :. All possible outcomes are:
- {(1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6)}
- :. Total number of possible outcomes = 36
- (i) Let *E* be the event that 5 does not come up either time.

Number of favourable outcomes = [36 - (5 + 6)] = 25

- $\therefore P(E) = \frac{25}{36}$
- (ii) Let N be the event that 5 will come up at least once, then number of favourable outcomes = 5 + 6 = 11
- $\therefore P(N) = \frac{11}{36}$
- **25.** (i) Given argument is not correct. Because, if two coins are tossed simultaneously then four outcomes are possible {HH, HT, TH, TT}. So total number of outcomes is 4.
- $\therefore$  The probability of each of these outcomes = 1/4.
- (ii) Correct. Because the two outcomes are possible. Total number of outcomes = 6 and odd numbers = 3 and even numbers = 3.

So, number of favourable outcomes = 3 (in both the cases even or odd ).

 $\therefore P(\text{getting an odd number}) = \frac{3}{6} = \frac{1}{2}$ 

#### EXERCISE - 15.2

- 1. Here, total number of possible outcomes =  $5 \times 5 = 25$
- (i) Outcomes for both customers visiting on same day are:

{(Tue., Tue.), (Wed., Wed.), (Thu., Thu.), (Fri., Fri.), (Sat., Sat.)}

Number of favourable outcomes = 5

- $\therefore$  Required probability =  $\frac{5}{25} = \frac{1}{5}$
- (ii) Outcomes for both the customers visiting on consecutive days are :

{(Tue., Wed.), (Wed., Thu.), (Thu., Fri.), (Fri., Sat.),

(Sat., Fri.), (Wed., Tue.), (Thu., Wed.), (Fri., Thu.)}

- $\Rightarrow$  Number of favourable outcomes = 8
- $\therefore$  Required probability =  $\frac{8}{25}$
- (iii) We have probability for both visiting on same day  $=\frac{1}{5}$
- ∴ Probability for both visiting on different days= 1 [Probability for both visiting on the same day]

$$=1-\frac{1}{5}=\frac{5-1}{5}=\frac{4}{5}$$

- $\therefore$  The required probability =  $\frac{4}{5}$ .
- 2. The completed table is as under:

1	2	2	3	3	6
2	3	3	4	4	7
3	4	4	5	5	8
3	4	4	5	5	8
4	5	5	6	6	9
4	5	5	6	6	9
7	8	8	9	9	12
	3 4 4	2 3 3 4 3 4 4 5 4 5	2 3 3 3 3 3 4 4 4 4 5 5 5 4 5 5	2 3 3 4 3 4 4 5 3 4 4 5 4 5 5 6 4 5 5 6	2 3 3 4 4 3 4 4 5 5 3 4 4 5 5 4 5 5 6 6 4 5 5 6 6

- ∴ Number of all possible outcomes = 36
- (i) For total score being even:

Favourable outcomes = 18

[: The even outcomes are: 2, 4, 4, 4, 4, 8, 4, 4, 8, 4, 6, 6, 4, 6, 6, 8, 8, 12]

- $\therefore \text{ The required probability } = \frac{18}{36} = \frac{1}{2}$
- (ii) For the total score being 6:

In list of scores, we have four 6's.

- :. Number of favourable outcomes = 4
- $\therefore$  Required probability =  $\frac{4}{36} = \frac{1}{9}$
- (iii) For the total score being at least 6:

The favourable scores are: 7, 8, 8, 6, 6, 9, 6, 6, 9, 7, 8, 8, 9, 9 and 12

- ⇒ Number of favourable outcomes = 15
- $\therefore \quad \text{Required probability} = \frac{15}{36} = \frac{5}{12}$
- 3. Let the number of blue balls in the bag be x.
- $\therefore$  Total number of balls = x + 5

Number of possible outcomes = (x + 5)

For a blue ball, number of favourable outcomes = x

 $\therefore$  Probability of drawing a blue ball =  $\frac{x}{x+5}$ 

Similarly, probability of drawing a red ball =  $\frac{5}{x+5}$ 

Now, we have 
$$\frac{x}{x+5} = 2\left[\frac{5}{x+5}\right]$$

$$\Rightarrow \quad \frac{x}{x+5} = \frac{10}{x+5} \Rightarrow x = 10$$

Thus, the required number of blue balls is 10.

- 4. : The total number of balls in the box = 12
- :. Total number of possible outcomes = 12

Case I: For drawing a black ball

Number of favourable outcomes = x

$$\therefore$$
 Probability of getting a black ball =  $\frac{x}{12}$ 

**Case II:** When 6 more black balls are added Now, the total number of balls = 12 + 6 = 18

- $\Rightarrow$  Total number of possible outcomes = 18 Now, the number of black balls = (x + 6).
- $\therefore$  Number of favourable outcomes = (x + 6)

$$\therefore \quad \text{Required probability} = \frac{x+6}{18}$$

According to the given condition,

$$\frac{x+6}{18} = 2\left(\frac{x}{12}\right)$$

$$\Rightarrow$$
 12 (x + 6) = 36x  $\Rightarrow$  12x + 72 = 36x

$$\Rightarrow 36x - 12x = 72 \Rightarrow 24x = 72 \Rightarrow x = \frac{72}{24} = 3$$

Thus, the required value of x is 3.

- 5. : There are 24 marbles in the jar.
- $\therefore$  Total number of possible outcomes = 24 Let there are x blue marbles in the jar.
- $\therefore$  Number of green marbles = 24 x
- $\Rightarrow$  Number of favourable outcomes = (24 x)
- $\therefore$  Required probability for drawing a green marble  $=\frac{24-x}{x}$

Now, according to the condition, we have  $\frac{24-x}{24} = \frac{2}{3}$ 

$$\Rightarrow$$
 3(24 - x) = 2 × 24  $\Rightarrow$  72 - 3x = 48

$$\Rightarrow 3x = 72 - 48 \Rightarrow 3x = 24 \Rightarrow x = \frac{24}{3} = 8$$

Thus, the required number of blue marbles is 8.

## MtG BEST SELLING BOOKS FOR CLASS 10







































