CHAPTER 7

Coordinate Geometry



SOLUTIONS

EXERCISE - 7.1

1. (i) Here, $x_1 = 2$, $y_1 = 3$ and $x_2 = 4$, $y_2 = 1$

:. The required distance

$$= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(4 - 2)^2 + (1 - 3)^2}$$
$$= \sqrt{2^2 + (-2)^2} = \sqrt{4 + 4} = \sqrt{8} = 2\sqrt{2} \text{ units}$$

(ii) Here, $x_1 = -5$, $y_1 = 7$ and $x_2 = -1$, $y_2 = 3$

:. The required distance

$$= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{[-1 - (-5)]^2 + (3 - 7)^2}$$
$$= \sqrt{(-1 + 5)^2 + (-4)^2} = \sqrt{16 + 16} = 4\sqrt{2} \text{ units}$$

(iii) Here $x_1 = a$, $y_1 = b$ and $x_2 = -a$, $y_2 = -b$

: The required distance

$$= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(-a - a)^2 + (-b - b)^2}$$

$$= \sqrt{(-2a)^2 + (-2b)^2} = \sqrt{4a^2 + 4b^2} = \sqrt{4(a^2 + b^2)}$$

$$= 2\sqrt{(a^2 + b^2)} \text{ units}$$

2. Part-I


Let the given points be A(0, 0) and B(36, 15).

Then,
$$AB = \sqrt{(36-0)^2 + (15-0)^2} = \sqrt{(36)^2 + (15)^2}$$

= $\sqrt{1296 + 225} = \sqrt{1521} = \sqrt{39^2} = 39$ units

Part-II

The given situation can be represented graphically as shown in the figure given below.



We have A(0, 0) and B(36, 15) as the positions of two towns.

Now,
$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

= $\sqrt{(36 - 0)^2 + (15 - 0)^2} = 39 \text{ km}.$

3. Let the given points be A(1, 5), B(2, 3) and C(-2, -11). Clearly, A, B and C will be collinear, if AB + BC = AC or AC + CB = AB or BA + AC = BC

Here,
$$AB = \sqrt{(2-1)^2 + (3-5)^2}$$

= $\sqrt{1^2 + (-2)^2} = \sqrt{1+4} = \sqrt{5} = 2.24$ units (Approx.)

$$BC = \sqrt{(-2-2)^2 + (-11-3)^2}$$

$$= \sqrt{(-4)^2 + (-14)^2} = \sqrt{16+196} = \sqrt{212} = 2\sqrt{53}$$
= 14.56 units (Approx.)

and,
$$AC = \sqrt{(-2-1)^2 + (-11-5)^2}$$

= $\sqrt{(-3)^2 + (-16)^2} = \sqrt{9 + 256} = \sqrt{265}$ units
= 16.28 units (Approx.)

Since, $AB + BC \neq AC$, $AC + CB \neq AB$ and $BA + AC \neq BC$ \therefore A, B and C are not collinear.

4. Let the given points be *A*(5, -2), *B*(6, 4) and *C*(7, -2).

Then,
$$AB = \sqrt{(6-5)^2 + [4-(-2)]^2}$$

= $\sqrt{(1)^2 + (6)^2} = \sqrt{1+36} = \sqrt{37}$ units

$$BC = \sqrt{(7-6)^2 + (-2-4)^2}$$
$$= \sqrt{(1)^2 + (-6)^2} = \sqrt{1+36} = \sqrt{37} \text{ units}$$

and
$$AC = \sqrt{(7-5)^2 + [-2-(-2)]^2}$$

= $\sqrt{(2)^2 + (0)^2} = \sqrt{4+0} = 2$ units

Since, AB = BC

 $\triangle ABC$ is an isosceles triangle.

5. The coordinates of given points are A(3, 4), B(6, 7), C(9, 4) and D(6, 1)

$$AB = \sqrt{(6-3)^2 + (7-4)^2}$$

$$= \sqrt{(3)^2 + (3)^2} = \sqrt{9+9} = \sqrt{18} = 3\sqrt{2} \text{ units}$$

$$BC = \sqrt{(9-6)^2 + (4-7)^2}$$

$$= \sqrt{3^2 + (-3)^2} = \sqrt{9+9} = \sqrt{18} = 3\sqrt{2} \text{ units}$$

$$CD = \sqrt{(6-9)^2 + (1-4)^2}$$

$$= \sqrt{(-3)^2 + (-3)^2} = \sqrt{9+9} = \sqrt{18} = 3\sqrt{2} \text{ units}$$

$$AD = \sqrt{(6-3)^2 + (1-4)^2}$$

$$= \sqrt{(3)^2 + (-3)^2} = \sqrt{9+9} = \sqrt{18} = 3\sqrt{2} \text{ units}$$

$$AC = \sqrt{(9-3)^2 + (4-4)^2} = \sqrt{(6)^2 + (0)^2} = 6 \text{ units}$$
and
$$BD = \sqrt{(6-6)^2 + (1-7)^2} = \sqrt{(0)^2 + (-6)^2} = 6 \text{ units}$$

Since, AB = BC = CD = AD *i.e.*, all the four sides are equal. and also, BD = AC *i.e.*, both the diagonals are also equal. $\therefore ABCD$ is a square.

Thus, Champa is correct.

6. (i) Let the given points be *A*(-1, -2), *B*(1, 0), *C*(-1, 2) and *D*(-3, 0).

Now,
$$AB = \sqrt{(1+1)^2 + (0+2)^2}$$

 $= \sqrt{(2)^2 + (2)^2} = \sqrt{4+4} = \sqrt{8}$ units
 $BC = \sqrt{(-1-1)^2 + (2-0)^2} = \sqrt{4+4} = \sqrt{8}$ units
 $CD = \sqrt{(-3+1)^2 + (0-2)^2} = \sqrt{4+4} = \sqrt{8}$ units
 $DA = \sqrt{(-1+3)^2 + (-2-0)^2} = \sqrt{4+4} = \sqrt{8}$ units
 $AC = \sqrt{(-1+1)^2 + (2+2)^2} = \sqrt{0+4^2} = 4$ units
 $BD = \sqrt{(-3-1)^2 + (0-0)^2} = \sqrt{(-4)^2} = 4$ units

Since, AB = BC = CD = DA *i.e.*, all the sides are equal, and also, AC = BD *i.e.*, the diagonals are also equal.

 \therefore ABCD is a square.

(ii) Let the given points be A(-3, 5), B(3, 1), C(0, 3) and D(-1, -4).

Now,
$$AB = \sqrt{[3 - (-3)]^2 + (1 - 5)^2}$$

 $= \sqrt{6^2 + (-4)^2} = \sqrt{36 + 16} = \sqrt{52} = 2\sqrt{13}$ units
 $BC = \sqrt{(0 - 3)^2 + (3 - 1)^2} = \sqrt{9 + 4} = \sqrt{13}$ units
 $CD = \sqrt{(-1 - 0)^2 + (-4 - 3)^2} = \sqrt{(-1)^2 + (-7)^2}$
 $= \sqrt{1 + 49} = \sqrt{50}$ units
 $DA = \sqrt{[-3 - (-1)]^2 + [5 - (-4)]^2} = \sqrt{(-2)^2 + (9)^2}$
 $= \sqrt{4 + 81} = \sqrt{85}$ units
 $AC = \sqrt{[0 - (-3)]^2 + (3 - 5)^2} = \sqrt{(3)^2 + (-2)^2}$
 $= \sqrt{9 + 4} = \sqrt{13}$ units
and $BD = \sqrt{(-1 - 3)^2 + (-4 - 1)^2} = \sqrt{(-4)^2 + (-5)^2}$
 $= \sqrt{16 + 25} = \sqrt{41}$ units

Here, we can see that $\left[\because \sqrt{13} + \sqrt{13} = 2\sqrt{13}\right]$ AC + BC = AB

 \Rightarrow A, B and C are collinear points. Hence, ABCD is not a quadrilateral.

(iii) Let the given points be A(4, 5), B(7, 6), C(4, 3) and D(1, 2).

Now,
$$AB = \sqrt{(7-4)^2 + (6-5)^2} = \sqrt{3^2 + 1^2} = \sqrt{10}$$
 units
 $BC = \sqrt{(4-7)^2 + (3-6)^2} = \sqrt{(-3)^2 + (-3)^2} = \sqrt{18}$ units
 $CD = \sqrt{(1-4)^2 + (2-3)^2} = \sqrt{(-3)^2 + (-1)^2} = \sqrt{10}$ units
 $DA = \sqrt{(4-1)^2 + (5-2)^2} = \sqrt{9+9} = \sqrt{18}$ units
 $AC = \sqrt{(4-4)^2 + (3-5)^2} = \sqrt{0 + (-2)^2} = 2$ units

and $BD = \sqrt{(1-7)^2 + (2-6)^2} = \sqrt{36+16} = \sqrt{52}$ units Since, AB = CD, BC = DA i.e., opposite sides of the given quadrilateral are equal, and also, $AC \neq BD$, i.e., diagonals are unequal.

∴ *ABCD* is a parallelogram.

7. We know that any point on x-axis is of the form (x, 0).

 \therefore Let the required point be P(x, 0).

Also, let the given points be A(2, -5) and B(-2, 9).

Now,
$$AP = \sqrt{(x-2)^2 + [0-(-5)]^2}$$

 $= \sqrt{(x-2)^2 + 5^2} = \sqrt{x^2 - 4x + 4 + 25} = \sqrt{x^2 - 4x + 29}$
and $BP = \sqrt{[x-(-2)]^2 + (0-9)^2}$
 $= \sqrt{(x+2)^2 + (-9)^2} = \sqrt{x^2 + 4x + 4 + 81} = \sqrt{x^2 + 4x + 85}$

Since, *A* and *B* are equidistant from *P*.

$$\therefore AP = BP$$

$$\Rightarrow \sqrt{x^2 - 4x + 29} = \sqrt{x^2 + 4x + 85}$$

$$\Rightarrow x^2 - 4x + 29 = x^2 + 4x + 85$$

$$\Rightarrow -8x = 56 \Rightarrow x = \frac{56}{9} = -7$$

 \therefore The required point is (-7, 0).

8. The given points are P(2, -3) and Q(10, y).

$$PQ = \sqrt{(10-2)^2 + [y - (-3)]^2}$$

$$= \sqrt{8^2 + (y+3)^2} = \sqrt{64 + y^2 + 6y + 9} = \sqrt{y^2 + 6y + 73}$$
But $PQ = 10$ [Given]

But
$$PQ = 10$$

∴ $\sqrt{y^2 + 6y + 73} = 10$

On squaring both sides, we get $y^2 + 6y + 73 = 100$

$$\Rightarrow y^2 + 6y - 27 = 0$$

$$\Rightarrow y^2 - 3y + 9y - 27 = 0 \Rightarrow (y - 3)(y + 9) = 0$$

$$\Rightarrow y = 3 \text{ or } y = -9$$

 \therefore The required values of *y* are 3 and -9.

9. Here,
$$QP = \sqrt{(5-0)^2 + (-3-1)^2} = \sqrt{5^2 + (-4)^2}$$

= $\sqrt{25+16} = \sqrt{41}$ units

and
$$QR = \sqrt{(x-0)^2 + (6-1)^2}$$

= $\sqrt{x^2 + 5^2} = \sqrt{x^2 + 25}$ units

$$\therefore QP = QR$$

$$\therefore \sqrt{41} = \sqrt{x^2 + 25}$$

On squaring both sides, we get $x^2 + 25 = 41$

$$\Rightarrow x^2 = 16 \Rightarrow x = \pm 4$$

Thus, the point R is (4, 6) or (-4, 6)

Now,
$$QR = \sqrt{[(\pm 4) - (0)]^2 + (6 - 1)^2} = \sqrt{16 + 25} = \sqrt{41}$$
 units
and $PR = \sqrt{(4 - 5)^2 + (6 + 3)^2}$ or $\sqrt{(-4 - 5)^2 + (6 + 3)^2}$
 $\Rightarrow PR = \sqrt{1 + 81}$ or $\sqrt{81 + 81}$

$$\Rightarrow$$
 $PR = \sqrt{82}$ units or $9\sqrt{2}$ units

10. Let A(x, y), B(3, 6) and C(-3, 4) be the given points. Now let, the point A(x, y) is equidistant from B(3, 6) and C(-3, 4).

Then, we get AB = AC

$$\Rightarrow \sqrt{(3-x)^2 + (6-y)^2} = \sqrt{(-3-x)^2 + (4-y)^2}$$

$$(3-x)^2 + (6-y)^2 = (-3-x)^2 + (4-y)^2$$

On squaring both sides, we get

$$(3-x)^2 + (6-y)^2 = (-3-x)^2 + (4-y)^2$$

$$\Rightarrow 9+x^2-6x+36+y^2-12y=9+x^2+6x+16+y^2-8y$$

$$\Rightarrow$$
 $-6x - 6x + 36 - 12y - 16 + 8y = 0$

$$\Rightarrow$$
 -12x - 4y + 20 = 0 \Rightarrow -3x - y + 5 = 0

 \Rightarrow 3x + y - 5 = 0, which is the required relation between x and y.

EXERCISE - 7.2

Let the required point be P(x, y). Here, the end points are (-1, 7) and (4, -3)

Ratio = $2:3 = m_1:m_2$

$$\therefore \quad x = \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}$$

$$=\frac{(2\times4)+3\times(-1)}{2+3}=\frac{8-3}{5}=\frac{5}{5}=1$$

and
$$y = \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} = \frac{2 \times (-3) + (3 \times 7)}{2 + 3}$$

$$=\frac{-6+21}{5}=\frac{15}{5}=3$$

Thus, the required point is (1, 3).

Let the points P and Q trisect AB

i.e., AP = PQ = QB

i.e., P divides AB in the ratio of 1:2 and Q divides AB in the ratio of 2:1.

Let the coordinates of P be (x, y)

$$\therefore x = \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2} = \frac{1(-2) + 2(4)}{1 + 2} = \frac{-2 + 8}{3} = 2 \text{ and}$$

$$y = \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} = \frac{1(-3) + 2 \times (-1)}{1 + 2} = \frac{-3 - 2}{3} = \frac{-5}{3}$$

 \therefore The required coordinates of *P* are $\left(2, \frac{-5}{2}\right)$.

Let the coordinates of
$$Q$$
 be (X, Y) .

$$\therefore X = \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2} = \frac{2(-2) + 1(4)}{2 + 1} = \frac{-4 + 4}{3} = 0$$

$$Y = \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} = \frac{2(-3) + 1(-1)}{2 + 1} = \frac{-6 - 1}{3} = \frac{-7}{3}$$

- The required coordinates of Q are $\left(0, \frac{-7}{2}\right)$.
- Let us consider A as origin, then AB is the x-axis and AD is the y-axis.

Now, the position of green flag-post is $\left(2, \frac{100}{4}\right)$ or (2, 25).

and, the position of red flag-post is $\left(8, \frac{100}{5}\right)$ or (8, 20).

Distance between both the flags

$$= \sqrt{(8-2)^2 + (20-25)^2}$$

$$= \sqrt{6^2 + (-5)^2} = \sqrt{36 + 25} = \sqrt{61} \,\mathrm{m}$$

Let the mid-point of the line segment joining the two flags be M(x, y).

$$\therefore x = \frac{2+8}{2} \text{ and } y = \frac{25+20}{2}$$

$$(2, 25) \qquad (x, y)$$

$$(2, 25)$$
 (x, y) $(8, 20)$

 \Rightarrow x = 5 and y = 22.5

Thus, the blue flag is on the 5th line at a distance 22.5 m

Given, points are A(-3, 10) and B(6, -8)

Let the point P(-1, 6) divides AB in the ratio k:1.

Using section formula, we have

$$(-1,6) = \left(\frac{6k-3}{k+1}, \frac{-8k+10}{k+1}\right) A(-3,10) B(6,-8)$$

$$\Rightarrow \frac{6k-3}{k+1} = -1$$
 and $\frac{-8k+10}{k+1} = 6$

⇒
$$6k - 3 = -k - 1$$
 and $-8k + 10 = 6k + 6$
⇒ $7k = 2$ and $14 = 4$

$$\Rightarrow$$
 7k = 2 and 14 k = 4

$$\Rightarrow k = \frac{2}{7}$$

Required ratio is $\frac{2}{7}$: 1 *i.e.*, 2:7.

The given points are A(1, -5) and B(-4, 5).

Let the required ratio be k: 1 and the required point be P(x, y).

Since the point *P* lies on *x*-axis,

Its y-coordinate is 0.

$$\therefore x = \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2} \text{ and } 0 = \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2}$$

$$\Rightarrow x = \frac{k(-4) + 1(1)}{k + 1} \text{ and } 0 = \frac{5k - 5}{k + 1}$$

$$\Rightarrow x = \frac{k(-4) + 1(1)}{k + 1}$$
 and $0 = \frac{5k - 5}{k + 1}$

$$\Rightarrow$$
 $x = \frac{-4k+1}{k+1}$ and $0 = \frac{5k-5}{k+1}$

$$\Rightarrow x(k+1) = -4k + 1 \text{ and } 5k - 5 = 0 \Rightarrow k = 1$$
$$\Rightarrow x(1+1) = -4 + 1$$

$$\Rightarrow x(1+1) = -4+1 \qquad [\because k=1]$$

$$\Rightarrow 2x = -3 \Rightarrow x = -\frac{3}{2}$$


 \therefore The required ratio is 1:1 and coordinates of *P* are $\left(\frac{-3}{2},0\right)$.

Let the given points are A(1, 2), B(4, y), C(x, 6)and D(3, 5).

Since, the diagonals of a parallelogram bisect each other.

The coordinates of *P*

$$X = \frac{x+1}{2} = \frac{3+4}{2}$$




$$\Rightarrow$$
 $x + 1 = 7 \Rightarrow x = 6$ and $Y = \frac{5 + y}{2} = \frac{6 + 2}{2}$

$$\Rightarrow$$
 5 + y = 8 \Rightarrow y = 3

The required values of x and y are 6 and 3 respectively.

Here, centre of the circle is O(2, -3).

Let the end points of the diameter be A(x, y) and B(1, 4).



The centre of a circle bisects the diameter.

$$\therefore 2 = \frac{x+1}{2} \Rightarrow x+1 = 4 \Rightarrow x = 3$$

And,
$$-3 = \frac{y+4}{2} \Rightarrow y+4 = -6 \Rightarrow y = -10$$

Hence, the coordinates of A are (3, -10).

8.
$$P(x, y)$$
 $A(-2, -2)$ 3 4 $B(2, -4)$

Here, the given points are A(-2, -2) and B(2, -4). Let the coordinates of P are (x, y).

Since, the point *P* lies on *AB* such that

$$AP = \frac{3}{7}AB \Rightarrow \frac{AP}{AB} = \frac{3}{7} \Rightarrow \frac{AB}{AP} = \frac{7}{3}$$

$$\Rightarrow \frac{AP + BP}{AP} = \frac{7}{3} \qquad (\because AB = AP + BP)$$

$$\Rightarrow 1 + \frac{BP}{AP} = \frac{3+4}{3} = 1 + \frac{4}{3} \Rightarrow \frac{BP}{AP} = \frac{4}{3}$$

$$\Rightarrow$$
 AP: PB = 3: 4 i.e., $P(x, y)$ divides AB in the ratio 3: 4.

$$\therefore x = \frac{3 \times 2 + 4 \times (-2)}{3 + 4} = \frac{6 - 8}{7} = \frac{-2}{7} \text{ and}$$
$$y = \frac{3 \times (-4) + 4 \times (-2)}{3 + 4} = \frac{-12 - 8}{7} = \frac{-20}{7}$$

Thus, the coordinates of P are $\left(-\frac{2}{7}, -\frac{20}{7}\right)$.

Here, the given points are A(-2, 2) and B(2, 8). Let P_1 , P_2 and P_3 divide AB in four equal parts.

$$A(-2, 2) P_1 P_2 P_3 B(2, 8)$$

Since, $AP_1 = P_1P_2 = P_2P_3 = P_3B$

 P_2 is the mid-point of AB

$$\therefore \quad \text{Coordinates of } P_2 \text{ are } \left(\frac{-2+2}{2}, \frac{2+8}{2} \right) = (0,5)$$

Again, P_1 is the mid-point of AP_2 .

 \therefore Coordinates of P_1 are

$$\left(\frac{-2+0}{2}, \frac{2+5}{2}\right) = \left(-1, \frac{7}{2}\right)$$

Also, P_3 is the mid-point of P_2B .

 \therefore Coordinates of P_3 are

$$\left(\frac{0+2}{2}, \frac{5+8}{2}\right) = \left(1, \frac{13}{2}\right)$$

Thus, the coordinates of P_1 , P_2 and P_3 are

$$\left(-1, \frac{7}{2}\right)$$
, $(0,5)$ and $\left(1, \frac{13}{2}\right)$ respectively.

10. Let the vertices of the given rhombus are A(3, 0), B(4, 5), C(-1, 4) and D(-2, -1).

 \therefore AC and BD are the diagonals of rhombus ABCD.

$$AC = \sqrt{(-1-3)^2 + (4-0)^2}$$

$$= \sqrt{(-4)^2 + (4)^2} = \sqrt{16+16} = 4\sqrt{2} \text{ units}$$

$$BD = \sqrt{(-2-4)^2 + (-1-5)^2}$$

$$= \sqrt{(-6)^2 + (-6)^2}$$

$$= \sqrt{36+36} = 6\sqrt{2} \text{ units}$$

$$A(3,0)$$

$$C(-1,4)$$

: Area of a rhombus

$$= \frac{1}{2} \times (\text{Product of diagonals})$$

$$= \frac{1}{2} \times AC \times BD$$

$$= \frac{1}{2} \times 4\sqrt{2} \times 6\sqrt{2} = 4 \times 6 = 24 \text{ square units.}$$

EXERCISE - 7.3

(i) Let the vertices of the triangles be A(2, 3), B(-1, 0)and C(2, -4). \therefore Area of a $\triangle ABC$

$$= \frac{1}{2} [2\{0 - (-4)\} + (-1)\{-4 - (3)\} + 2\{3 - 0\}]$$

$$= \frac{1}{2} [2(0 + 4) + (-1)(-4 - 3) + 2(3)]$$

$$= \frac{1}{2} [8 + 7 + 6] = \frac{1}{2} [21] = \frac{21}{2} \text{ sq.units}$$

(ii) Let the vertices of the triangles be A(-5, -1), B(3, -5)and C(5, 2).

Area of a
$$\triangle ABC$$

$$= \frac{1}{2} [-5\{-5-2\} + 3\{2-(-1)\} + 5\{-1-(-5)\}]$$

$$= \frac{1}{2} [-5\{-7\} + 3\{2+1\} + 5\{-1+5\}]$$

$$= \frac{1}{2} [35+9+20] = \frac{1}{2} \times 64 = 32 \text{ sq.units}$$

- The given three points will be collinear if the area of triangle formed by them is zero.
- (i) Let A(7, -2), B(5, 1) and C(3, k) be the vertices of a triangle.
- The given points will be collinear, if ar $(\Delta ABC) = 0$

$$\Rightarrow \frac{1}{2} [7(1-k) + 5(k+2) + 3(-2-1)] = 0$$

 $\Rightarrow 7 - 7k + 5k + 10 - 9 = 0$

$$\Rightarrow$$
 8 - 2k = 0 \Rightarrow 2k = 8 \Rightarrow k = $\frac{8}{2}$ = 4

(ii) Let A(8, 1), B(k, -4) and C(2, -5) be the vertices of a triangle.

The given points will be collinear, if $ar(\Delta ABC) = 0$

$$\Rightarrow \frac{1}{2}[8(-4+5)+k(-5-1)+2(1+4)]=0$$

\Rightarrow 8-6k+10=0 \Rightarrow 6k = 18 \Rightarrow k = 3.

3. Let the vertices of the triangle be A(0, -1), B(2, 1) and C(0, 3).


Let D, E and F be the mid-points of the sides BC, CA and AB respectively.

Then, coordinates of D are

$$\left(\frac{2+0}{2}, \frac{1+3}{2}\right) = (1,2)$$

Coordinates of E are

$$\left(\frac{0+0}{2}, \frac{3+(-1)}{2}\right) = (0,1)$$



Coordinates of *F* are
$$\left(\frac{2+0}{2}, \frac{1+(-1)}{2}\right) = (1, 0)$$

Now,
$$\operatorname{ar}(\Delta ABC) = \frac{1}{2}[0(1-3) + 2\{3 - (-1)\} + 0(-1-1)]$$

= $\frac{1}{2}[0 + 8 + 0] = \frac{1}{2} \times 8 = 4 \text{ sq.units}$

Now,
$$\operatorname{ar}(\Delta DEF) = \frac{1}{2}[1(1-0)+0(0-2)+1(2-1)]$$

= $\frac{1}{2}[1+0+1] = \frac{1}{2} \times 2 = 1 \text{ sq.unit}$

$$\therefore \frac{\operatorname{ar}(\Delta DEF)}{\operatorname{ar}(\Delta ABC)} = \frac{1}{4}; \quad \therefore \operatorname{ar}(\Delta DEF) : \operatorname{ar}(\Delta ABC) = 1 : 4$$

4. Let A(-4, -2), B(-3, -5), C(3, -2) and D(2, 3) be the vertices of the quadrilateral.

Let us join diagonal BD.

Now, $ar(\Delta ABD)$

$$= \frac{1}{2}[(-4)\{-5-3\} + (-3)\{3-(-2)\} + 2\{(-2)-(-5)\}]$$


$$= \frac{1}{2}[(-4)(-8) + (-3)(5) + 2(-2+5)]$$

$$= \frac{1}{2}[32 + (-15) + 6] = \frac{1}{2}[23] = \frac{23}{2} \text{ sq.units}$$
Also, $\operatorname{ar}(\triangle CBD)$

$$= \frac{1}{2} [3(-5-3) + (-3)(3-(-2)) + 2((-2) - (-5))]$$

$$= \frac{1}{2}[3(-8) + (-3)(5) + 2(3)]$$
$$= \frac{1}{2}[-24 - 15 + 6]$$

$$=\frac{1}{2}[-33]=-\frac{33}{2}$$



$$\therefore \quad \text{ar } (\Delta CBD) = \frac{33}{2} \text{ sq., units}$$

[: Area of triangle cannot be negative] ARCD = ar(AARD) + ar(ACRD)

Since, $ar(quad. ABCD) = ar(\Delta ABD) + ar(\Delta CBD)$

∴ ar(quad.
$$ABCD$$
) = $\left(\frac{23}{2} + \frac{33}{2}\right)$ sq. units
= $\frac{56}{2}$ sq.units = 28 sq.units

5. Here, the vertices of the triangle are A(4, -6), B(3, -2) and C(5, 2).

Let *D* be the mid-point of *BC*.

The coordinates of the mid- A(4, -6) point of D are $\left\{\frac{3+5}{2}, \frac{-2+2}{2}\right\} = (4, 0)$ Since AD divides the triangle ABC into two parts i.e., $\triangle ABD$ and $\triangle ADC$, Now, $\operatorname{ar}(\triangle ABD)$ $= \frac{1}{2}[4\{(-2)-0\}+3(0+6)+4(-6+2)]$

$$= \frac{1}{2} [4\{(-2) - 0\} + 3(0 + 6) + 4(-6 + 2)]$$
$$= \frac{1}{2} [(-8) + 18 + (-16)] = \frac{1}{2} (-6) = -3$$

$$\therefore \quad \text{ar}(\Delta ABD) = 3 \text{ sq. units} \qquad \dots (i)$$

[: Area of triangle cannot be negative]

Also,
$$\operatorname{ar}(\Delta ADC) = \frac{1}{2}[4(0-2) + 4(2+6) + 5(-6-0)]$$

= $\frac{1}{2}[-8 + 32 - 30] = \frac{1}{2}[-6] = -3$

$$\therefore$$
 ar($\triangle ADC$) = 3 sq. units ...(ii)

[: Area of triangle cannot be negative]

From (i) and (ii), we have $ar(\Delta ABD) = ar(\Delta ADC)$

Hence, a median divides the triangle into two triangles of equal areas.

EXERCISE - 7.4

1. Let the point C divides the line segment joining the points A(2, -2) and B(3, 7) in the ratio k : 1. Using section formula, we have

Coordinates of C are
$$\left(\frac{3k+2}{k+1}, \frac{7k-2}{k+1}\right)$$

Since, the point *C* lies on the given line 2x + y - 4 = 0.

.. We have
$$2\left(\frac{3k+2}{k+1}\right) + \left(\frac{7k-2}{k+1}\right) - 4 = 0$$

 $\Rightarrow 2(3k+2) + (7k-2) - 4(k+1) = 0$

$$\Rightarrow 2(3k+2) + (7k-2) - 4(k+1) = 0$$

$$\Rightarrow 6k+4+7k-2-4k-4=0$$

$$\Rightarrow 6k + 4 + 7k - 2 - 4k - 4 =$$

$$\Rightarrow 9k - 2 = 0 \Rightarrow k = \frac{2}{9}$$

 \therefore The required ratio is $\frac{2}{9}$: 1 *i.e.*, 2:9.

2. Let the given points be A(x, y), B(1, 2) and C(7, 0). Given, the points A, B and C will be collinear.

.. Area of
$$\triangle ABC = 0 \Rightarrow \frac{1}{2} [x(2-0) + 1(0-y) + 7(y-2)] = 0$$

 $\Rightarrow 2x - y + 7y - 14 = 0$
 $\Rightarrow 2x + 6y - 14 = 0 \Rightarrow x + 3y - 7 = 0$, which is the

 \Rightarrow 2x + 6y - 14 - 0 \Rightarrow x + 3y - 7 - 0, which is the required relation between x and y.

3. Let P(x, y) be the centre of the circle and the circle is passing through the points A(6, -6), B(3, -7) and C(3, 3). ∴ AP = BP = CP

Taking AP = BP, we have $AP^2 = BP^2$

$$\Rightarrow$$
 $(x-6)^2 + (y+6)^2 = (x-3)^2 + (y+7)^2$

$$\Rightarrow x^2 - 12x + 36 + y^2 + 12y + 36 = x^2 - 6x + 9 + y^2 + 14y + 49$$

$$\Rightarrow -12x + 6x + 12y - 14y + 72 - 58 = 0$$

$$\Rightarrow$$
 $-6x - 2y + 14 = 0$

$$\Rightarrow 3x + y - 7 = 0 \qquad ... (i)$$
Taking $BP = CP$, we have
$$BP^2 = CP^2$$

$$\Rightarrow (x - 3)^2 + (y + 7)^2 = (x - 3)^2 + (y - 3)^2$$

$$\Rightarrow x^2 - 6x + 9 + y^2 + 14y + 49 = x^2 - 6x + 9 + y^2 - 6y + 9$$

$$\Rightarrow 14y + 6y + 58 - 18 = 0 \Rightarrow 20y + 40 = 0$$

$$\Rightarrow y = \frac{-40}{20} = -2 \qquad ... (ii)$$

From (i) and (ii), we get


$$3x - 2 - 7 = 0$$
 \Rightarrow $3x = 9 \Rightarrow x = 3$

x = 3 and y = -2

Hence, the required centre is (3, -2).

Let us have a square ABCD such that A(-1, 2) and C(3, 2) are the opposite vertices.

Let B(x, y) be an unknown vertex. Since, all sides of a square are equal.



$$\therefore AB = CB \Rightarrow AB^2 = CB^2$$

$$\Rightarrow (x+1)^2 + (y-2)^2 = (x-3)^2 + (y-2)^2$$

$$\Rightarrow x^2 + 1 + 2x + y^2 + 4 - 4y = x^2 + 9 - 6x + y^2 + 4 - 4y$$

$$\Rightarrow 2x + 1 = -6x + 9 \Rightarrow 8x = 8 \Rightarrow x = 1 \qquad \dots (i)$$

Since, each angle of a square = 90°

ABC is a right angled triangle.

Using Pythagoras theorem, we have

$$AB^2 + CB^2 = AC^2$$

$$AB^{2} + CB^{2} = AC^{2}$$

$$\Rightarrow [(x+1)^{2} + (y-2)^{2}] + [(x-3)^{2} + (y-2)^{2}]$$

$$= [(3+1)^{2} + (2-2)^{2}]$$

$$= [(3+1)^{2} + (2-2)^{2}]$$

$$\Rightarrow 2x^2 + 2y^2 + 2x - 4y - 6x - 4y + 1 + 4 + 9 + 4 = 16$$

$$\Rightarrow 2x^2 + 2y^2 - 4x - 8y + 2 = 0$$

$$\Rightarrow x^2 + y^2 - 2x - 4y + 1 = 0$$
 ... (ii

Substituting the value of *x* from (i) into (ii), we have

$$1 + y^2 - 2 - 4y + 1 = 0$$

$$\Rightarrow y^2 - 4y = 0 \Rightarrow y(y - 4) = 0$$

$$\Rightarrow$$
 $y = 0$ or $y = 4$

Hence, the two required other vertices are (1, 0) and (1, 4).

- (i) By taking A as the origin and AD and AB as the coordinate axes. We have P(4, 6), Q(3, 2) and R(6, 5) as the vertices of ΔPQR .
- (ii) By taking C as the origin and CB and CD as the coordinate axes, then the vertices of $\triangle PQR$ are P(-12, -2), Q(-13, -6) and R(-10, -3).

Now area of ΔPQR [when P(4, 6), Q(3, 2) and R(6, 5) are

$$= \frac{1}{2}[4(2-5) + 3(5-6) + 6(6-2)]$$
$$= \frac{1}{2}[-12 - 3 + 24] = \frac{9}{2} \text{ sq. units}$$

Area of $\triangle PQR$ [when P(-12, -2), Q(-13, -6) and R(-10, -3)are the vertices

$$= \frac{1}{2}[-12(-6+3) + (-13)(-3+2) + (-10)(-2+6)]$$

$$= \frac{1}{2}[-12(-3) + (-13)(-1) + (-10)4]$$

$$= \frac{1}{2}[36+13-40] = \frac{9}{2} \text{ sq.units}$$

Thus, in both cases, the area of $\triangle PQR$ is the same.

6. We have,
$$\frac{AD}{AB} = \frac{1}{4}$$

$$\Rightarrow \frac{AB}{AD} = \frac{4}{1} \Rightarrow \frac{AD + DB}{AD} = \frac{4}{1}$$

$$\Rightarrow \frac{AD}{AD} + \frac{DB}{AD} = \frac{4}{1}$$

$$\Rightarrow 1 + \frac{DB}{AD} = 1 + \frac{3}{1} \Rightarrow \frac{DB}{AD} = \frac{3}{1}$$

$$\Rightarrow AD \cdot DB = 1 \cdot 3$$

Thus, the point D divides AB in the ratio 1:3.

The coordinates of *D* are

$$\left[\frac{(1\times1)+(3\times4)}{1+3}, \frac{(1\times5)+(3\times6)}{1+3}\right]$$

$$=\left[\frac{1+12}{4}, \frac{5+18}{4}\right] = \left[\frac{13}{4}, \frac{23}{4}\right]$$

Similarly, AE : EC = 1 : 3 i.e., E divides AC in the ratio 1 : 3.

Coordinates of E are

$$\left[\frac{(1\times7) + (3\times4)}{1+3}, \frac{1\times2 + 3\times6}{1+3}\right]$$
$$= \left[\frac{7+12}{4}, \frac{2+18}{4}\right] = \left[\frac{19}{4}, 5\right]$$

$$= \frac{1}{2} \left[4 \left(\frac{23}{4} - 5 \right) + \frac{13}{4} (5 - 6) + \frac{19}{4} \left(6 - \frac{23}{4} \right) \right]$$

$$= \frac{1}{2} \left[(23 - 20) + \frac{13}{4} (-1) + \frac{19}{4} \left(\frac{24 - 23}{4} \right) \right]$$


$$= \frac{1}{2} \left[3 - \frac{13}{4} + \frac{19}{16} \right] = \frac{1}{2} \left[\frac{48 - 52 + 19}{16} \right] = \frac{15}{32} \text{ sq. units}$$

Area of
$$\triangle ABC = \frac{1}{2}[4(5-2)+1(2-6)+7(6-5)]$$

$$= \frac{1}{2}[(4\times3)+1(-4)+7\times1]$$

$$= \frac{1}{2}[12+(-4)+7] = \frac{1}{2}(15) = \frac{15}{2} \text{ sq.units}$$
Now, $\frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle ABC)} = \frac{\frac{15}{32}}{\frac{15}{2}} = \frac{15}{32} \times \frac{2}{15} = \frac{1}{16}$

- \Rightarrow ar($\triangle ADE$): ar($\triangle ABC$) = 1:16.
- We have the vertices of $\triangle ABC$ as A(4, 2), B(6, 5) and C(1, 4).



- Since *AD* is a median
- D is the mid-point of BC.

$$\therefore$$
 Coordinates of *D* are $\left(\frac{6+1}{2}, \frac{5+4}{2}\right) = \left(\frac{7}{2}, \frac{9}{2}\right)$

(ii) Since AP : PD = 2 : 1 *i.e.,* P divides AD in the ratio 2 : 1.

 \therefore Coordinates of P are

$$\left(\frac{2\left(\frac{7}{2}\right) + (1 \times 4)}{2+1}, \frac{2\left(\frac{9}{2}\right) + 1 \times 2}{2+1}\right) = \left(\frac{7+4}{3}, \frac{9+2}{3}\right) = \left(\frac{11}{3}, \frac{11}{3}\right)$$

(iii) Since, BE is the median.

$$\therefore$$
 Coordinates of E are $\left(\frac{4+1}{2}, \frac{2+4}{2}\right) = \left(\frac{5}{2}, 3\right)$

 $BQ: QE = 2: 1 \Rightarrow$ The point Q divides BE in the ratio 2: 1.

 \therefore Coordinates of Q are

$$\left(\frac{2\left(\frac{5}{2}\right)+1\times 6}{2+1}, \frac{(2\times 3)+(1\times 5)}{2+1}\right)$$
$$=\left(\frac{5+6}{3}, \frac{6+5}{3}\right) = \left(\frac{11}{3}, \frac{11}{3}\right)$$

Since, CF is the median.

$$\therefore$$
 Coordinates of F are $\left(\frac{4+6}{2}, \frac{2+5}{2}\right) = \left(5, \frac{7}{2}\right)$

 \Rightarrow The point *R* divides *CF* in the ratio 2 : 1

So, Coordinates of *R* are

$$\left(\frac{2 \times 5 + 1 \times 1}{2 + 1}, \frac{2 \times \frac{7}{2} + 1 \times 4}{2 + 1}\right)$$


$$= \left(\frac{10 + 1}{3}, \frac{7 + 4}{3}\right) = \left(\frac{11}{3}, \frac{11}{3}\right)$$

(iv) We observe that P, Q and R represent the same point.

(v) Here, we have $A(x_1, y_1)$, $B(x_2, y_2)$ and $C(x_3, y_3)$ are the vertices of $\triangle ABC$. Let AD, BE and CF are its medians.

 \therefore D, E and F are the mid-points of BC, CA and AB

respectively.



 $A(x_1, y_1)$

We know, the centroid is a point on a median, dividing it in the ratio 2:1.

Considering the median AD, coordinates of D are

$$\left[\frac{x_2+x_3}{2}, \frac{y_2+y_3}{2}\right]$$

Let *G* be the centroid.

.. Coordinates of the centroid are

$$\left[\frac{(1 \times x_1) + 2\left(\frac{x_2 + x_3}{2}\right)}{1 + 2}, \frac{(1 \times y_1) + 2\left(\frac{y_2 + y_3}{2}\right)}{1 + 2}\right]$$
$$= \left[\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}\right]$$

Similarly, considering the other medians we find that in each the coordinates of *G* are


$$\left(\frac{x_1+x_2+x_3}{3}, \frac{y_1+y_2+y_3}{3}\right)$$

i.e., The coordinates of the centroid are

$$\left(\frac{x_1+x_2+x_3}{3}, \frac{y_1+y_2+y_3}{3}\right)$$

8. We have a rectangle whose vertices are A(-1, -1), B(-1, 4), C(5, 4) and D(5, -1).

7



 \therefore *P* is mid-point of *AB*

$$\therefore$$
 Coordinates of *P* are $\left(\frac{-1-1}{2}, \frac{-1+4}{2}\right) = \left(-1, \frac{3}{2}\right)$

Similarly, coordinates of Q are $\left(\frac{-1+5}{2}, \frac{4+4}{2}\right) = (2,4)$

Coordinates of R are
$$\left(\frac{5+5}{2}, \frac{-1+4}{2}\right) = \left(5, \frac{3}{2}\right)$$

Coordinates of S are
$$\left(\frac{-1+5}{2}, \frac{-1-1}{2}\right) = (2,-1)$$

Now,
$$PQ = \sqrt{(2+1)^2 + \left(4 - \frac{3}{2}\right)^2} = \sqrt{9 + \frac{25}{4}} = \frac{\sqrt{61}}{2}$$
 units

$$QR = \sqrt{(2-5)^2 + \left(4 - \frac{3}{2}\right)^2} = \sqrt{9 + \frac{25}{4}} = \frac{\sqrt{61}}{2}$$
 units

$$RS = \sqrt{(2-5)^2 + \left(-1 + \left(-\frac{3}{2}\right)^2} = \sqrt{9 + \frac{25}{4}} = \frac{\sqrt{61}}{2} \text{ units}$$

$$SP = \sqrt{(2+1)^2 + \left(-1 - \frac{3}{2}\right)^2} = \sqrt{9 + \frac{25}{4}} = \frac{\sqrt{61}}{2}$$
 units

$$PR = \sqrt{(5+1)^2 + \left(\frac{3}{2} - \frac{3}{2}\right)^2} = \sqrt{6^2 + 0} = 6 \text{ units}$$

$$QS = \sqrt{(2-2)^2 + (4+1)^2} = \sqrt{0+5^2} = 5$$
 units

We see that PQ = QR = RS = SP *i.e.*, all sides of quadrilateral PQRS are equal.

:. It can be a square or a rhombus.

But its diagonals are not equal.

i.e.,
$$PR \neq QS$$

 \therefore *PQRS* is a rhombus.

MtG BEST SELLING BOOKS FOR CLASS 10



