



## TRY YOURSELF

## SOLUTIONS

1. Let  $n = 5q + 1$ , where  $q$  is a positive integer  
Squaring both sides, we get  
 $n^2 = (5q + 1)^2 = 25q^2 + 10q + 1$   
 $= 5(5q^2 + 2q) + 1 = 5m + 1$ ,  
where  $m = 5q^2 + 2q$  is an integer  
Hence, the square of any positive integer of the form  $5q + 1$  is of the same form.

2. On dividing  $n$  by 3, let  $q$  be the quotient and  $r$  be the remainder.

Then, by Euclid's division lemma,

$$n = 3q + r, \text{ where } 0 \leq r < 3$$

$$\Rightarrow n = 3q \text{ or } n = 3q + 1 \text{ or } n = 3q + 2$$

**Case I :** If  $n = 3q$ , which is divisible by 3  
but  $(n + 1)$  and  $(n + 2)$  are not divisible by 3.

So, in this case, only  $n$  is divisible by 3.

**Case II :** If  $n = 3q + 1$ , then  $n + 2 = 3q + 3$

$$= 3(q + 1) \text{ which is divisible by 3}$$

but  $n$  and  $(n + 1)$  are not divisible by 3.

So, in this case, only  $(n + 2)$  is divisible by 3.

**Case III :** If  $n = 3q + 2$ , then  $n + 1 = 3q + 3$

$$= 3(q + 1) \text{ which is divisible by 3}$$

but  $n$  and  $(n + 2)$  are not divisible by 3.

So, in this case, only  $(n + 1)$  is divisible by 3.

Thus, one and only one out of  $n$ ,  $(n + 1)$  and  $(n + 2)$  is divisible by 3.

3. Here,  $250 > 30$

$\therefore$  Applying Euclid's division lemma, we get

$$250 = 30 \times 8 + 10$$

Since, remainder,  $10 \neq 0$

$\therefore$  Applying Euclid's division lemma to 30 and 10, we get

$$30 = 10 \times 3 + 0$$

Since, remainder is 0, when divisor is 10

$\therefore$  By Euclid's division algorithm,

$$\text{HCF}(250, 30) = 10$$

4. The required number of soaps in each box is HCF of 612 and 342.

By Euclid's division algorithm, we have

$$612 = 342 \times 1 + 270,$$

$$342 = 270 \times 1 + 72,$$

$$270 = 72 \times 3 + 54,$$

$$72 = 54 \times 1 + 18,$$

$$54 = 18 \times 3 + 0$$

Here, remainder is 0, when divisor is 18

$$\therefore \text{HCF}(612, 342) \text{ is } 18$$

So, the trader can pack 18 soaps in each box.

5. Given numbers are 1305, 1365 and 1530

$$\therefore 1530 > 1365 > 1305$$

$\therefore$  By applying Euclid's division lemma to 1530 and 1365, we get

$$1530 = 1365 \times 1 + 165,$$

$$1365 = 165 \times 8 + 45,$$

$$165 = 45 \times 3 + 30,$$

$$45 = 30 \times 1 + 15, 30 = 15 \times 2 + 0$$

Since, remainder is 0 when divisor is 15

$$\therefore \text{HCF}(1530, 1365) \text{ is } 15$$

Now, applying Euclid's division lemma to 1305 and 15, we get


$$1305 = 15 \times 87 + 0$$

Since, remainder is 0, when divisor is 15

$$\therefore \text{HCF}(1305, 15) \text{ is } 15$$


Hence, HCF of 1305, 1365 and 1530 is 15.

6. (i) Using factor tree method, we have



$$\therefore 4095 = 3 \times 3 \times 5 \times 7 \times 13 = 3^2 \times 5 \times 7 \times 13$$

(ii) Using factor tree method, we have



$$\therefore 1001 = 7 \times 11 \times 13$$

7. We have,  $9 \times 13 \times 17 + 17$

$= 17(9 \times 13 + 1) = 17(117 + 1) = 17 \times 118$ , which is not a prime number because it has 17 as a factor other than 1 and the number itself.

$\therefore 9 \times 13 \times 17 + 17$  is a composite number.

Also, we have  $5 \times 6 \times 7 \times 8 \times 9 + 7 = 7(5 \times 6 \times 8 \times 9 + 1)$ , which is again not a prime number because it has 7 as a factor other than 1 and the number itself.

$\therefore 5 \times 6 \times 7 \times 8 \times 9 + 7$  is a composite number.

8. If any number ends with the digit 0 or 5, it is always divisible by 5.

If  $12^n$  ends with the digit zero or five, it must be divisible by 5.

This is possible only if prime factorisation of  $12^n$  contains the prime number 5.

$$\text{Now, } 12 = 2 \times 2 \times 3 = 2^2 \times 3$$

$$\Rightarrow 12^n = (2^2 \times 3)^n = 2^{2n} \times 3^n$$

Since, there is no term containing 5.

Hence, there is no value of  $n$  for which  $12^n$  ends with the digit zero or five.

9. The prime factorisation of 144, 180 and 192 is,

$$144 = 2 \times 2 \times 2 \times 2 \times 3 \times 3 = 2^4 \times 3^2$$

$$180 = 2 \times 2 \times 3 \times 3 \times 5 = 2^2 \times 3^2 \times 5$$

$$192 = 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 3 = 2^6 \times 3$$

$$\therefore \text{HCF}(144, 180, 192) = 2^2 \times 3 = 12$$

$$\text{and LCM}(144, 180, 192) = 2^6 \times 3^2 \times 5 = 2880$$

10. Since, the books are to be distributed equally among the students of section A or section B.

So, number of books must be a multiple of 32 as well as 36

$\therefore$  Required number of books is the LCM of 32 and 36

Prime factorisation of 32 and 36 is

$$32 = 2 \times 2 \times 2 \times 2 \times 2 = 2^5$$

$$36 = 2 \times 2 \times 3 \times 3 = 2^2 \times 3^2$$

$$\therefore \text{LCM of } 32 \text{ and } 36 = 2^5 \times 3^2 = 288$$

Hence, required number of books is 288.

11. Given, HCF ( $a, b$ ) = 11

Product of  $a$  and  $b$  = 7623

$$\therefore \text{LCM}(a, b) = \frac{\text{Product of } a \text{ and } b}{\text{HCF}(a, b)}$$

$$\Rightarrow \text{LCM}(a, b) = \frac{7623}{11} = 693$$

12. Given HCF (2520, 6600) = 120

$$\text{LCM}(2520, 6600) = 252 \times k$$

$$\text{Now, HCF}(2520, 6600) \times \text{LCM}(2520, 6600) = 2520 \times 6600$$

$$\Rightarrow (120) \times (252 \times k) = 2520 \times 6600$$

$$\Rightarrow k = \frac{2520 \times 6600}{252 \times 120} = 550$$

13. We know that HCF ( $a, b$ )  $\times$  LCM ( $a, b$ ) = product of  $a$  and  $b$

$$\Rightarrow 12 \times \text{LCM}(a, b) = 1152$$

$$\Rightarrow \text{LCM}(a, b) = \frac{1152}{12} = 96$$

14. Given,  $x = p^2 q^3$ ,  $y = p^3 q$ , where  $p$  and  $q$  are primes.

$$\text{LCM}(x, y) = p^3 q^3$$

$$\text{HCF}(x, y) = p^2 q$$

$$\text{Now, LCM}(x, y) = p^3 q^3 = p q^2 p^2 q = p q^2 \times \text{HCF}(x, y)$$

$\therefore$  LCM is a multiple of HCF.

15. Let us assume that  $\sqrt{3}$  is rational

So, we can find integers  $a$  and  $b$  ( $b \neq 0$  and  $a, b$  are co-prime) such that

$$\sqrt{3} = \frac{a}{b} \Rightarrow \sqrt{3}b = a$$

$$\Rightarrow 3b^2 = a^2 \quad [\text{Squaring both sides}]$$

$$\therefore 3 \text{ divides } a^2 \Rightarrow 3 \text{ divides } a \quad \dots(\text{ii})$$

So, we can write  $a = 3m$ , where  $m$  is an integer

Putting  $a = 3m$  in (i), we get

$$3b^2 = 9m^2 \Rightarrow b^2 = 3m^2$$

$$\therefore 3 \text{ divides } b^2 \Rightarrow 3 \text{ divides } b \quad \dots(\text{iii})$$

From (ii) and (iii), 3 is a common factor of  $a$  and  $b$ , which contradicts the fact that  $a$  and  $b$  are co-prime.

$\therefore$  Our assumption that  $\sqrt{3}$  is rational is wrong.

Hence,  $\sqrt{3}$  is irrational.

16. Let us assume that  $3 + \sqrt{2}$  is rational.

So, we can find two integers  $a$  and  $b$  ( $b \neq 0$  and  $a, b$  are co-prime)

$$\text{such that } 3 + \sqrt{2} = \frac{a}{b}, \Rightarrow \sqrt{2} = \frac{a}{b} - 3 = \frac{a - 3b}{b}$$

$$\text{Here, } \frac{a - 3b}{b} \text{ is rational} \quad [\because a \text{ and } b \text{ are integers}]$$

$\Rightarrow \sqrt{2}$  is rational, which contradicts the fact that  $\sqrt{2}$  is irrational

$\therefore$  Our supposition is wrong.

Hence,  $3 + \sqrt{2}$  is irrational.

17. We have a rational number 23.3408, whose decimal expansion terminates.

$$\begin{aligned} \text{Now, } 23.3408 &= \frac{233408}{10000} = \frac{233408}{10^4} = \frac{2^6 \times 7 \times 521}{2^4 \times 5^4} \\ &= \frac{2^2 \times 7 \times 521}{5^4} = \frac{14588}{2^0 \times 5^4} \end{aligned}$$

Thus, 23.3408 can be expressed as  $\frac{14588}{2^0 \times 5^4}$ , where

numerator and denominator are co-prime and denominator is of the form  $2^m \times 5^n$ , where  $m, n$  are non-negative integers.

18. Here, denominator,

$$2^2 \times 15^3 = 2^2 \times (5 \times 3)^3 = 2^2 \times 5^3 \times 3^3$$

$$\therefore \frac{129}{2^2 \times 5^3 \times 3^3} \text{ will be a non-terminating repeating}$$

decimal expansion because it contains 3 as a factor in its denominator.

19. As, we need only 10 as denominator, so we multiply

$$\frac{1}{3} \text{ with } \frac{3}{10}$$

$$\therefore \frac{1}{3} \times \frac{3}{10} = \frac{1}{10} = 0.1$$

